

University of Groningen

Ankylosing spondylitis disease activity score is related to NSAID use, especially in patients treated with TNF-alpha inhibitors

Carbo, Marlies J G; Spoorenberg, Anneke; Maas, Fiona; Brouwer, Elisabeth; Bos, Reinhard; Bootsma, Hendrika; van der Veer, Eveline; Wink, Freke; Arends, Suzanne

Published in:
 PLoS ONE

DOI:
 [10.1371/journal.pone.0196281](https://doi.org/10.1371/journal.pone.0196281)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
 Publisher's PDF, also known as Version of record

Publication date:
 2018

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Carbo, M. J. G., Spoorenberg, A., Maas, F., Brouwer, E., Bos, R., Bootsma, H., van der Veer, E., Wink, F., & Arends, S. (2018). Ankylosing spondylitis disease activity score is related to NSAID use, especially in patients treated with TNF-alpha inhibitors. *PLoS ONE*, 13(4), [e0196281].
 <https://doi.org/10.1371/journal.pone.0196281>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.


RESEARCH ARTICLE

Ankylosing spondylitis disease activity score is related to NSAID use, especially in patients treated with TNF- α inhibitors

Marlies J. G. Carbo^{1*}, Anneke Spoorenberg^{1,2}, Fiona Maas¹, Elisabeth Brouwer¹, Reinhard Bos², Hendrika Bootsma¹, Eveline van der Veer³, Freke Wink², Suzanne Arends^{1,2}

1 Rheumatology and Clinical Immunology, University Medical Center Groningen, University of Groningen, Groningen, the Netherlands, **2** Rheumatology, Medical Center Leeuwarden, Leeuwarden, the Netherlands, **3** Laboratory Medicine, University Medical Center Groningen, University of Groningen, Groningen, the Netherlands

* m.j.g.carbo@umcg.nl



Abstract

OPEN ACCESS

Citation: Carbo MJG, Spoorenberg A, Maas F, Brouwer E, Bos R, Bootsma H, et al. (2018) Ankylosing spondylitis disease activity score is related to NSAID use, especially in patients treated with TNF- α inhibitors. PLoS ONE 13(4): e0196281. <https://doi.org/10.1371/journal.pone.0196281>

Editor: Tatsuo Kanda, Nihon University School of Medicine, JAPAN

Received: December 20, 2017

Accepted: April 10, 2018

Published: April 24, 2018

Copyright: © 2018 Carbo et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Data are available from the University of Groningen—UMCG Institutional Data Access for researchers who meet the criteria for access to confidential data. The local ethics committees of the Medical Center Leeuwarden (MCL) and the University Medical Center Groningen (UMCG) will maintain the ethical restrictions of the data. The Data Protection Officer of the UMCG will maintain the legal restrictions and appropriate codes of conduct. Permission is required prior to access. Data requests can be sent

Background

Non-steroidal anti-inflammatory drugs (NSAIDs) are regarded as the cornerstone of conventional treatment for AS. However little is known about concomitant NSAID use during treatment (with TNF- α inhibitors) in daily clinical practice.

Methods and findings

Consecutive patients from the GLAS cohort were included. NSAID use and ASAS-NSAID index were evaluated at group level and at individual patient level during 52 weeks of follow-up. Analyses were stratified for treatment regimen. Generalized estimating equations (GEE) was used to evaluate NSAID use in relation to assessments of disease activity over time. In patients starting TNF- α inhibitors ($n = 254$), 79% used NSAIDs at baseline and this proportion decreased significantly to 38% at 52 weeks. ASAS-NSAID index also decreased significantly from median 65 to 0. In patients on conventional treatment ($n = 139$), 74% used NSAIDs at baseline with median ASAS-NSAID index of 50 and this remained stable during follow-up. At each follow-up visit, approximately half of the patients changed their type or dose of NSAIDs. GEE analysis over time showed that NSAID use was associated with AS disease activity score ($p < 0.05$). This relation was more pronounced in patients treated with TNF- α inhibitors compared to conventional treatment ($B = 0.825$ vs. $B = 0.250$).

Conclusions

In this observational cohort of established AS patients, there was no difference in baseline NSAID use between patients with and without indication for TNF- α inhibitors. NSAID use decreased significantly after starting TNF- α inhibitors. During conventional treatment,

to Research Data Office University of Groningen:
researchdata@rug.nl.

Funding: The GLAS cohort was supported by an unrestricted grant from Pfizer. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have the following interests: MJG Carbo: None declared; Spoorenberg: Grant/research support from Abbvie, Pfizer, UCB, consultant for Abbvie, Pfizer, MSD, UCB, and Novartis; F. Maas: none declared; E. Brouwer: Grant/research support from Pfizer; R. Bos: Grant/research support from Abbvie, Pfizer, consultant for Pfizer and BMS; H. Bootsma: None declared; E van der Veer: None declared; F.R. Wink: consultant for Abbvie; S. Arends: Grant/research support from Pfizer. This does not alter our adherence to PLOS ONE policies on sharing data and materials.

NSAID use remained stable at group level. However, NSAID use changed frequently at individual patient level and was significantly associated with disease activity.

Introduction

Ankylosing spondylitis (AS) is a chronic inflammatory rheumatic disease that primarily affects the axial skeleton. Non-steroidal anti-inflammatory drugs (NSAIDs) are regarded as the cornerstone of conventional treatment for AS.[1] NSAIDs have shown good reduction of symptoms in 60–80% of the patients.[2] A recent network meta-analysis of randomized controlled trials (RCTs), reporting on the efficacy of different NSAIDs in AS, showed that the majority of available NSAIDs reduce total pain score significantly compared to placebo up to 12 weeks.[3] Besides the positive effect on pain, NSAIDs also reduce the level of acute-phase reactants in the blood of AS patients.[4] Furthermore, a decrease in signal intensity of bone marrow edema of the sacroiliac (SI) joints on MRI was seen after 6 weeks of full dose NSAIDs in newly diagnosed patients with axial spondyloarthritis (SpA).[5] During treatment, disadvantages of NSAIDs such as possible cardiovascular and gastrointestinal side effects should be taken into account, especially in patients with comorbidity and comedication (e.g. anticoagulants).[6,7]

In AS, there is only limited data available comparing the efficacy of continued use of NSAIDs with on demand use. A single RCT studied symptom control and safety of continued versus on demand use of celecoxib and ketoprofen during 2 years of follow-up as a secondary outcome. No differences were found between the groups.[7] A recent Cochrane review on both traditional and COX-2 selective NSAIDs in AS found no difference in harms between NSAIDs and placebo during 12 weeks of follow-up.[8] Conflicting results were published about the effect of continued versus on demand use of NSAIDs on spinal radiographic progression in AS.[9,10]

For over a decade, tumor necrosis factor-alpha (TNF- α) inhibitors are available for AS patients with active disease who have insufficient response to conventional treatment including NSAIDs. TNF- α inhibitors have shown to reduce the clinical signs and symptoms as well as serum levels of CRP and axial inflammation detected on MRI in AS patients with active disease.[11] A head to head comparison between NSAIDs and TNF- α inhibitors on efficacy in treatment naïve patients has never been performed. Additionally, little is known about the additional efficacy of concomitant NSAID use to the treatment of biologicals in AS. Therefore, it can be questioned whether AS patients should be advised to stop or continue their NSAIDs during TNF- α inhibitor use.[5]

So far, studies on concomitant NSAID use to the treatment of biologicals were only performed in patients with early active axiale SpA.[12–15] A recent RCT showed that patients reached partial remission more frequently during treatment with infliximab combined with naproxen than during naproxen treatment alone.[16] Another recent RCT showed the NSAID sparing effect of etanercept treatment. Patients were able to reduce their NSAID intake by more than half during 8 weeks of etanercept treatment.[15] In the observational DESIR cohort, patients were included presenting with inflammatory back pain, symptom duration between ≥ 3 months and 3 years and symptoms suggestive of spondyloarthritis according to the local investigator. However, these patients did not necessarily fulfill the modified New York criteria for AS.[13] Patients receiving TNF- α inhibitors from this cohort were matched to patients on conventional treatment using propensity scores. After 2 years of follow-up, in both treatment groups the proportion of patients using NSAIDs decreased as well as the Assessment of Spondyloarthritis international Society (ASAS)-NSAID index, which is based on the specific

NSAID, dosage and frequency. These results were significantly more pronounced in the group receiving TNF- α inhibitors.[15] Based on these results, it was concluded that most patients can reduce their intake of NSAIDs after starting TNF- α inhibitors. However, not all patients receiving TNF- α inhibitors were able to stop or decrease their NSAID intake.

The aim of the present study was to evaluate NSAID use over time and to investigate if NSAID use was related to disease activity in patients with established AS from a large observational cohort study starting TNF- α inhibitors or receiving conventional treatment during a follow-up period of 52 weeks.

Methods

Patients

The GLAS cohort was approved by the local ethics committees of the MCL and UMCG and all patients provided written informed consent according to the Declaration of Helsinki. The present analysis was performed in the Groningen Leeuwarden Ankylosing Spondylitis (GLAS) cohort. The GLAS cohort is an ongoing prospective longitudinal observational cohort study with standardized follow-up. Since November 2004, consecutive AS outpatients who started TNF- α inhibitors from the University Medical Center Groningen (UMCG) and the Medical Center Leeuwarden (MCL) were included.[17] In 2009, this inclusion was extended to all consecutive axial SpA outpatients, irrespective of treatment regimen. All patients were 18 years or older and fulfilled the modified New York criteria for AS (>90%) or the ASAS criteria for axial SpA including MRI. All patients included before June 2013 were included in this analysis. Patients were excluded if they started TNF- α inhibitors prior to inclusion in the GLAS cohort or had an indication but did not start with TNF- α inhibitors due to non-medical reasons, such as the patient's own preference (Fig 1).

Treatment

NSAIDs were prescribed based on expert opinion. TNF- α inhibitors were started according to the ASAS consensus statement.[18] The choice between available TNF- α inhibitors was based on the judgment of the treating rheumatologist and/or the specific preference of the patient. [17] After 3 months of TNF- α inhibitor treatment, patients with low disease activity were advised to take NSAIDs on demand. There was no difference in treatment advice between males and females.

Follow-up and clinical assessments

Visits were scheduled at baseline (before start of TNF- α inhibitors) and after 6, 12, 24 and 52 weeks for the TNF- α inhibitor group. The conventional treatment group had scheduled visits at baseline, 24 and 52 weeks. At every visit, disease activity was assessed using the AS Disease Activity Score (ASDAS), Bath AS Disease Activity Index (BASDAI), and serum CRP level. [19–20]

At every visit, NSAID use (yes/no) and type of NSAID was recorded. Dosage and frequency were assessed retrospectively from clinical records to calculate the ASAS-NSAID index. This index takes the specific NSAID and dosage into account as well as the percentage of days of NSAID intake during a period of interest.[1,21] A score of 100 is equal to a full dose of NSAIDs all days of the week. For situations in which the exact amount of NSAID use was unclear, a frequency of 2 days a week was used for occasional NSAID use (1 to 3 days a week) and 0.5 days a week for very rare NSAID use (<1 day a week). To determine the changes in NSAID use, the




Fig 1. Flowchart of inclusion.

<https://doi.org/10.1371/journal.pone.0196281.g001>

type and dosage of NSAIDs was assessed at each visit to investigate starting, switching, dose escalation, dose reduction or complete stop of NSAIDs.

Statistics

Results were expressed as number of patients (%) and mean \pm SD or median (interquartile range; IQR), for categorical and normally or non-normally distributed data, respectively. Analyses were stratified for treatment regimen: patients starting TNF- α inhibitors and patients on conventional treatment. Chi-square or Fisher's Exact tests, Independent Samples t-tests, and Mann-Whitney U-tests were used to compare baseline characteristics. Linear and logistic generalized estimating equations (GEE) were used to analyze NSAID use over time. GEE is a technique for longitudinal data analysis, which makes use of all available data and allows unequal numbers of repeated measurements.[22] The 'exchangeable' correlation structure was used to correct for within patient correlation. If there was no normal distribution of residuals, parameters were log-transformed before being entered into the equation.

GEE was also applied to analyze the relation between NSAID use and disease activity over time, where disease activity was defined as the dependent variable. NSAID use was analyzed using 4 parameters: NSAID use (yes versus no), ASAS-NSAID index (continuous scale), low (index ≥ 10 versus < 10), and high use (index ≥ 90 versus < 90). To exclude potential bias due to TNF- α inhibitors discontinuation, analyses were also performed in patients using TNF- α inhibitors $\geq 80\%$ of the follow up time. To exclude the influence of the initial positive effect of TNF- α inhibitors on the association between NSAID use and disease activity, analyses were

also performed for 12 to 52 weeks of follow-up. P-values <0.05 were considered statistically significant. Statistical analysis was performed with IBM SPSS Statistics 22 (SPSS, Chicago, IL, USA).

Results

Of the 393 included AS patients, 67% were male, mean age was 44 ± 13 years, median symptom duration 15 years (IQR 8–24), and 79% were HLA-B27 positive. Additionally, 254 (66%) patients started TNF- α inhibitors due to active disease (etanercept (58%), adalimumab (26%), or infliximab (15%)). Of the 254 patients who started TNF- α inhibitors, 217 (85%) used this treatment more than 80% of the follow-up time. The remaining 139 (34%) patients received conventional treatment. Patient characteristics were comparable between both groups, except higher disease activity, more often peripheral arthritis, and worse physical functioning in patients starting TNF- α inhibitors.(Table 1)

NSAID use at group level

At baseline, 77% of patients used 12 different NSAIDs, mostly diclofenac (16%), naproxen (16%) and piroxicam (7%).(S1 Table) Baseline NSAID use was comparable between patients who started TNF- α inhibitors and patients on conventional treatment (79% vs. 74%). In patients who started TNF- α inhibitors, this proportion decreased significantly during follow-

Table 1. Baseline characteristics of the AS study population.

	Total n = 393	TNF- α inhibitors n = 254	Conventional treatment n = 139	P-value*
Patient characteristics				
Age (yrs)	44 \pm 13	43 \pm 12	44 \pm 14	0.451
Male gender	261 (66%)	173 (68%)	88 (64%)	0.253
HLA-B27+	294 (79%)	197 (81%)	97 (75%)	0.094
Duration of symptoms (yrs)	15 (8–24)	15 (9–24)	14 (7–24)	0.393
Time since diagnosis (yrs)	6 (1–16)	7 (1–24)	5 (1–16)	0.693
History of extra-spinal manifestations				
Peripheral arthritis	102 (27%)	76 (31%)	26 (20%)	0.015
IBD	36 (10%)	23 (9%)	13 (10%)	0.495
Uveitis	108 (28%)	73 (29%)	35 (27%)	0.334
Psoriasis	28 (7%)	15 (6%)	13 (10%)	0.125
Disease status				
BASDAI (range 0 to 10)	5.3 \pm 2.2	6.1 \pm 1.7	3.9 \pm 2.2	<0.001
ASDAS	3.3 \pm 1.1	3.8 \pm 0.8	2.4 \pm 0.9	<0.001
CRP (mg/l)	8 (8–17)	13 (5–22)	3 (2–7)	<0.001
ESR (mm/h)	14 (7–28)	21 (10–35)	9 (4–13)	<0.001
Peripheral arthritis	50 (13%)	43 (17%)	7 (5%)	<0.001
BASFI (range 0 to 10)	4.8 \pm 2.5	5.7 \pm 2.1	3.2 \pm 2.3	<0.001
Concomitant medication use				
Current NSAID use	289 (77%)	194 (79%)	95 (74%)	0.131
Current DMARD use	57 (15%)	47 (19%)	10 (7%)	0.002

*p-value: TNF- α inhibitors vs. conventional treatment. Values are number of patients (percentage), mean \pm SD or median (IQR). HLA-B27+, human leukocyte antigen B27 positive; IBD, inflammatory bowel disease; BASDAI, Bath Ankylosing Spondylitis Disease Activity Index; ASDAS, Ankylosing Spondylitis Disease Activity Score; CRP, C-reactive protein; ESR, erythrocyte sedimentation rate; BASFI, Bath Ankylosing Spondylitis Functional Index; DMARD, disease-modifying antirheumatic drug

<https://doi.org/10.1371/journal.pone.0196281.t001>

Table 2. Changes in NSAID use at group level.

	Baseline	6 weeks	12 weeks	24 weeks	52 weeks	p- value
TNF-α inhibitors	n = 245	n = 214	n = 223	n = 223	n = 217	
NSAID use (Y/N)	79%	57%	47%	41%	38%	<0.001
ASAS-NSAID index	65 (23–100)	25 (0–60)	0 (0–50)	0 (0–50)	0 (0–50)	<0.001
NSAID use low	24%	47%	56%	76%	77%	<0.001
NSAID use high	44%	21%	15%	16%	14%	<0.001
Conventional treatment	n = 129			n = 109	n = 112	
NSAID use (Y/N)	74%			80%	82%	0.183
ASAS-NSAID index	50 (0–100)			50 (5–100)	50 (7–100)	0.382
NSAID use low	44%			31%	28%	0.448
NSAID use high	36%			37%	38%	0.721

Values are percentage of patients or median (IQR).

<https://doi.org/10.1371/journal.pone.0196281.t002>

up to 38% at 52 weeks. When analyzing 2 different subgroups, low on demand and high NSAID use (defined by ASAS-NSAID as index resp. <10 and ≥ 90), the largest decrease in NSAID use was found during the first 6 weeks, followed by a slower gradual decrease up to 52 weeks.(Table 2) In the conventional treatment group, the proportion of patients using NSAIDs remained stable over time (82% at 52 weeks).(Table 2)

Patients starting TNF- α inhibitors also showed a rapid and significant decrease in ASAS-NSAID index during the first 6 to 12 weeks and this reduction remained stable during all follow-up visits. In patients on conventional treatment, the total ASAS-NSAID index and subgroups did not change over time.(Table 2) These patterns were found in both males and females (data not shown). GEE analysis showed no significant interaction between gender and ASAS-NSAID index over time.

In total, 22 patients (6%) never used NSAIDs, of which 18 patients in the TNF- α inhibitor group and 4 patients in the conventional treatment group. In the TNF- α inhibitor group, 9 patients had known contraindications for NSAID use (gastrointestinal (n = 4), renal (n = 4) and a proven allergy (n = 1)). In the conventional treatment group, there were no known contraindications.

NSAID use at individual patient level

Assessing the variability in NSAID use at individual patient level revealed that 46 to 67% of the patients treated with TNF- α inhibitors remained on a stable dose at each follow-up visit. Furthermore, 20 to 46% of the patients reduced their dose or stopped NSAIDs at a follow-up visit. Additionally, only 8 to 15% of the patients increased their dose, started or switched NSAIDs at a follow-up visit.(Table 3) Of the patients using NSAIDs at baseline, 48% had stopped their NSAIDs 52 weeks after starting TNF- α inhibitors.

Although NSAID use at group level was stable over time in the conventional treatment group, there were many changes at individual patient level. At each follow-up visit, approximately half of the patients changed their NSAID use: 26 to 34% of the patients showed dose escalation, start or switch of NSAIDs and 21 to 26% dose reduction or stop of NSAIDs. (Table 3) Only 7% used NSAIDs at baseline and not at 52 weeks, where 13% did not use NSAIDs at baseline but had started at 52 weeks of follow up.

NSAID use in relation to disease activity

Disease activity reduced significantly after starting TNF- α inhibitors and remained low and stable during follow-up. In the conventional treatment group, disease activity was low and

Table 3. Changes in NSAID use at individual patient level.

	Baseline	6 weeks	12 weeks	24 weeks	52 weeks
TNF-α Inhibitors					
NSAID use at baseline	79%				
Dose escalation, start or switch		8%	11%	13%	15%
Stable dose		46%	59%	67%	61%
Dose reduction or stop		46%	30%	20%	24%
Conventional treatment					
NSAID use at baseline	74%				
Dose escalation, start or switch				34%	26%
Stable dose				46%	49%
Dose reduction or stop				21%	26%

Patients who used no NSAIDs during follow-up were excluded from this table: 11% in TNF- α Inhibitor group and 6% in the conventional treatment group.

<https://doi.org/10.1371/journal.pone.0196281.t003>

stable at all visits.(Fig 2) At 52 weeks, disease activity was comparable between both groups, BASDAI mean 3.4 vs. 3.7, ASDAS mean 2.2 vs. 2.3, and CRP median 3 vs. 3 respectively.

GEE analysis over time showed that NSAID use was significantly associated with disease activity. In the TNF- α inhibitor group, a significant association of all 4 NSAID parameters (NSAID use (yes/no), ASAS-NSAID index, low on demand use and high use) with ASDAS was found. GEE revealed that higher NSAID use was related to higher ASDAS and vice versa. (Table 4) Comparable results were found for BASDAI and CRP.(S2 and S3 Tables)

The association between NSAID use and ASDAS remained significant in patients who used TNF- α inhibitors more than 80% of the follow-up time and when analyzing only 12 to 52 weeks of follow-up to exclude the initial effect of TNF- α inhibitors, although the regression coefficients were lower in these last analyses. (S4 Table)

In the conventional treatment group, a significant but less prominent association of 3 NSAID parameters NSAID use (yes/no), low on demand use and high use) with ASDAS was found. (Table 4) BASDAI was only significantly associated with low on demand NSAID use. For CRP, no significant associations with NSAID use were found.

Discussion

In the present prospective observational cohort study, we evaluated NSAID use in relation to disease activity during 52 weeks of follow-up in established AS patients treated with and without TNF- α inhibitors in daily clinical practice. This is the first study assessing the difference in NSAID use at group level and at individual patient level in established AS.

In the TNF- α inhibitor group, a rapid and significant decrease of concomitant NSAID use was seen. NSAID use at group level remained low during the entire follow-up period. Although patients in our cohort were advised to take their NSAID on demand after 3 months of TNF- α inhibitor treatment, many patients reduced or stopped NSAIDs on their own initiative even earlier. The reason for this is most likely the positive effect on symptoms of TNF- α inhibitors. Our finding of rapid NSAID reduction after 6–12 weeks of TNF- α inhibitor treatment is in line with a previous RCT in early axial SpA which showed the NSAID sparing effect of 8 weeks etanercept treatment.[15] Additionally, early axial SpA patients of the DESIR cohort showed a gradual decline in NSAID use after starting TNF- α inhibitors during 2 years of follow-up.[14]

In our conventional treatment group, NSAID use remained stable over 52 weeks of follow-up, whereas in the conventional care group of the DESIR cohort a significant decrease in




Fig 2. Disease activity over time in AS patients. ASDAS over time in patients starting TNF- α inhibitors (A) and receiving conventional treatment (B); BASDAI over time in patients starting TNF- α inhibitors (C) and receiving conventional treatment (D); CRP over time in patients starting TNF- α inhibitors (E) and receiving conventional treatment (F). * p <0.05 compared to baseline. Box-and-whisker plots: boxes indicate medians with interquartile ranges; + indicate means; whiskers indicate 5–95 percentile; • indicate outliers.

<https://doi.org/10.1371/journal.pone.0196281.g002>

NSAID use over 2 years was shown.[14] The GLAS cohort is an established AS cohort in contrast to the DESIR cohort, which includes patients with recent-onset (<3 years) inflammatory back pain suggestive of axial SpA.[14] Furthermore, DESIR used propensity scores to match the conventional treatment group to the TNF- α inhibitor group. This may explain the differences in the course of NSAID use between the conventional treatment groups of both cohorts. In concordance with our results, in the DESIR cohort the decrease in NSAID use of patients

Table 4. Association between ASDAS and NSAID use over time in AS patients.

	B (95% CI)	P-value	Interval	n
TNF-α Inhibitors				
NSAID use Yes	0.825 (0.664–0.985)	<0.001	1074	251
ASAS-NSAID index	0.009 (0.007–0.012)	<0.001	1073	251
NSAID use low	-0.831 (-0.672– -0.990)	<0.001	1073	251
NSAID use high	0.855 (0.682–1.028)	<0.001	1073	251
Conventional treatment				
NSAID use Yes	0.250 (0.006–0.493)	0.045	315	131
ASAS-NSAID index	0.002 (0.000–0.005)	0.059	314	131
NSAID use low	- 0.223 (-0.425–0.022)	0.030	314	131
NSAID use high	0.269 (0.038–0.501)	0.023	314	131

<https://doi.org/10.1371/journal.pone.0196281.t004>

on conventional treatment was also less pronounced than in patients starting TNF- α inhibitor group.[14]

In our GLAS cohort, frequent inter-individual changes in NSAID use were observed at each follow-up visit in both treatment groups. Interestingly, GEE analysis over time showed that NSAID use was significantly associated with disease activity. In the TNF- α inhibitor group, this relation was most pronounced and significant for all disease activity measures (i.e. ASDAS, BASDAI and CRP). Results remained significant when including patients who used TNF- α inhibitors more than 80% of the follow-up time and when analyzing 12 to 52 weeks of follow-up to exclude the influence of the initial positive effect of TNF- α inhibitors on the GEE analyses.

In the conventional treatment group, the association between NSAID use and disease activity was less pronounced and only found for ASDAS. This can be explained by less variation in disease activity in this treatment group, e.g. the large majority of patients had normal CRP levels, but it can also be related to lower power in GEE analysis because of the lower number of included patients and less frequent follow-up visits.

Since this is an observational cohort study in AS patients, it reflects NSAID use in daily clinical practice. No sub-analyses were performed on specific NSAIDs, e.g. traditional and COX-2 NSAIDs, due to small numbers. However, a recent meta-analysis showed no clear superiority in efficacy of any specific NSAID.[3] Unfortunately, systematic data on the side effects of NSAIDs were missing in our study, which potentially could have explained why patients reduced or stopped their NSAID intake. Despite this limitation, a positive association between NSAID use and disease activity was still found.

In summary, this is the first study investigating NSAID use in an established AS cohort on TNF- α inhibitors or conventional treatment. At baseline, we found no differences in NSAID use between AS patients with and without indication for TNF- α inhibitors. After starting TNF- α inhibitors, NSAID use decreased rapidly and remained low during follow-up. In patients receiving conventional treatment, NSAID use remained stable at group level over time. Conversely, at individual patient level, both the type and dose of NSAIDs changed frequently irrespective of treatment regimen. Therefore, stable NSAID use at group level does not necessarily reflect stable NSAID use at individual patient level. This should be taken into account in future evaluations and research on NSAID use in axial SpA. Most importantly, this is the first prospective cohort study of established AS patients that showed NSAID use was significantly associated with assessments of disease activity, which was most pronounced for patients treated with TNF- α inhibitors.

Supporting information

S1 Table. Specific NSAIDs used at baseline.
(DOCX)

S2 Table. Association between BASDAI and NSAID use over time in AS patients.
*Subgroup analysis of patients who used TNF- α inhibitors $\geq 80\%$ of the follow up time. ** Analysis for 12 to 52 weeks of follow-up (excluding baseline and 6 weeks).
(DOCX)

S3 Table. Association between CRP and NSAID use over time in AS patients. *Subgroup analysis of patients who used TNF- α inhibitors $\geq 80\%$ of the follow up time. ** Analysis for 12 to 52 weeks of follow-up (excluding baseline and 6 weeks).
(DOCX)

S4 Table. Association between ASDAS and NSAID use over time in AS patients. *Subgroup analysis of patients who used TNF- α inhibitors $\geq 80\%$ of the follow up time. ** Analysis for 12 to 52 weeks of follow-up (excluding baseline and 6 weeks).
(DOCX)

Acknowledgments

The authors would like to thank all patients who participated in the GLAS cohort. Furthermore, the authors wish to acknowledge Dr. P.M. Houtman, Mrs. W. Gerlofs, Mrs. S. Katerbarg, Mrs. A. Krol, Mrs. R. Rumph, and Mrs. B. Burmania for their contribution to clinical data collection.

Author Contributions

Conceptualization: Marlies J. G. Carbo, Anneke Spoorenberg, Fiona Maas, Elisabeth Brouwer, Reinhard Bos, Hendrika Bootsma, Eveline van der Veer, Freke Wink, Suzanne Arends.

Data curation: Marlies J. G. Carbo, Fiona Maas, Eveline van der Veer, Freke Wink, Suzanne Arends.

Formal analysis: Marlies J. G. Carbo, Anneke Spoorenberg, Fiona Maas, Suzanne Arends.

Funding acquisition: Anneke Spoorenberg, Suzanne Arends.

Investigation: Marlies J. G. Carbo, Anneke Spoorenberg, Hendrika Bootsma, Suzanne Arends.

Methodology: Marlies J. G. Carbo, Anneke Spoorenberg, Fiona Maas, Elisabeth Brouwer, Reinhard Bos, Hendrika Bootsma, Freke Wink, Suzanne Arends.

Project administration: Anneke Spoorenberg, Fiona Maas, Suzanne Arends.

Resources: Anneke Spoorenberg, Fiona Maas, Suzanne Arends.

Supervision: Anneke Spoorenberg, Elisabeth Brouwer, Suzanne Arends.

Validation: Suzanne Arends.

Visualization: Marlies J. G. Carbo, Anneke Spoorenberg, Fiona Maas, Elisabeth Brouwer, Reinhard Bos, Hendrika Bootsma, Eveline van der Veer, Suzanne Arends.

Writing – original draft: Marlies J. G. Carbo, Anneke Spoorenberg, Fiona Maas, Suzanne Arends.

Writing – review & editing: Marlies J. G. Carbo, Anneke Spoorenberg, Fiona Maas, Elisabeth Brouwer, Reinhard Bos, Hendrika Bootsma, Eveline van der Veer, Freke Wink, Suzanne Arends.

References

1. Dougados M, Baeten D. Spondyloarthritis. *Lancet* 2011 Jun 18; 377(9783):2127–2137. [https://doi.org/10.1016/S0140-6736\(11\)60071-8](https://doi.org/10.1016/S0140-6736(11)60071-8) PMID: 21684383
2. Poddubnyy D, van der Heijde D. Therapeutic controversies in spondyloarthritis: nonsteroidal anti-inflammatory drugs. *Rheum Dis Clin North Am* 2012 Aug; 38(3):601–611. <https://doi.org/10.1016/j.rdc.2012.08.005> PMID: 23083758
3. Wang R, Dasgupta A, Ward MM. Comparative efficacy of non-steroidal anti-inflammatory drugs in ankylosing spondylitis: a Bayesian network meta-analysis of clinical trials. *Ann Rheum Dis* 2015 Aug 6.
4. Benhamou M, Gossec L, Dougados M. Clinical relevance of C-reactive protein in ankylosing spondylitis and evaluation of the NSAIDs/coxibs' treatment effect on C-reactive protein. *Rheumatology (Oxford)* 2010 Mar; 49(3):536–541.
5. Varkas G, Jans L, Cypers H, Van Praet L, Carron P, Elewaut D, et al. Brief Report: Six-Week Treatment of Axial Spondyloarthritis Patients With an Optimal Dose of Nonsteroidal Antiinflammatory Drugs: Early Response to Treatment in Signal Intensity on Magnetic Resonance Imaging of the Sacroiliac Joints. *Arthritis Rheumatol* 2016 Mar; 68(3):672–678. <https://doi.org/10.1002/art.39474> PMID: 26473982
6. Henriksson K, From J, Stratelis G. Patient-reported adherence to coprescribed proton pump inhibitor gastroprotection in osteoarthritis, rheumatoid arthritis, and ankylosing spondylitis patients using nonsteroidal anti-inflammatory drugs. *Patient Prefer Adherence* 2014 Nov 18; 8:1611–1617. <https://doi.org/10.2147/PPA.S70651> PMID: 25429206
7. Guellec D, Nocturne G, Tatar Z, Pham T, Sellam J, Cantagrel A, et al. Should non-steroidal anti-inflammatory drugs be used continuously in ankylosing spondylitis? *Joint Bone Spine* 2014 Jul; 81(4):308–312. <https://doi.org/10.1016/j.jbspin.2014.01.003> PMID: 24589253
8. Kroon FP, van der Burg LR, Ramiro S, Landewe RB, Buchbinder R, Falzon L, et al. Nonsteroidal Antiinflammatory Drugs for Axial Spondyloarthritis: A Cochrane Review. *J Rheumatol* 2016 Mar; 43(3):607–617. <https://doi.org/10.3899/jrheum.150721> PMID: 26834216
9. Wanders A, Heijde D, Landewe R, Behier JM, Calin A, Olivieri I, et al. Nonsteroidal antiinflammatory drugs reduce radiographic progression in patients with ankylosing spondylitis: a randomized clinical trial. *Arthritis Rheum* 2005 Jun; 52(6):1756–1765. <https://doi.org/10.1002/art.21054> PMID: 15934081
10. Kroon F, Landewe R, Dougados M, van der Heijde D. Continuous NSAID use reverts the effects of inflammation on radiographic progression in patients with ankylosing spondylitis. *Ann Rheum Dis* 2012 Oct; 71(10):1623–1629. <https://doi.org/10.1136/annrheumdis-2012-201370> PMID: 22532639
11. Braun J, Baraliakos X, Heldmann F, Kiltz U. Tumor necrosis factor alpha antagonists in the treatment of axial spondyloarthritis. *Expert Opin Investig Drugs* 2014 May; 23(5):647–659. <https://doi.org/10.1517/13543784.2014.899351> PMID: 24654630
12. Varkas G, Van den Bosch F. NSAIDs in axial spondyloarthritis: to be continued. . . ? *Ann Rheum Dis* 2016 Feb 26.
13. Dougados M, d'Agostinob M, Benessianoc J, Berenbaumd F, Brebane F, Claudepierre P, et al. Study design and baseline characteristics of the 708 recruited patients. *Joint Bone Spine*. December 2011, 78 (6), Pages 598–603 <https://doi.org/10.1016/j.jbspin.2011.01.013> PMID: 21458351
14. Molto A, Granger B, Wendling D, Breban M, Dougados M, Gossec L. Brief Report: Nonsteroidal Antiinflammatory Drug-Sparing Effect of Tumor Necrosis Factor Inhibitors in Early Axial Spondyloarthritis: Results From the DESIR Cohort. *Arthritis Rheumatol* 2015 Sep; 67(9):2363–2368. <https://doi.org/10.1002/art.39208> PMID: 26109532
15. Dougados M, Wood E, Combe B, Schaefferbeke T, Miceli-Richard C, Berenbaum F, et al. Evaluation of the nonsteroidal anti-inflammatory drug-sparing effect of etanercept in axial spondyloarthritis: results of the multicenter, randomized, double-blind, placebo-controlled SPARSE study. *Arthritis Res Ther* 2014 Nov 27; 16(6):481–481-0481-5.
16. Sieper J, Listing J, Poddubnyy D, Song IH, Hermann KG, Callhoff J, et al. Effect of continuous versus on-demand treatment of ankylosing spondylitis with diclofenac over 2 years on radiographic progression of the spine: results from a randomised multicentre trial (ENRADAS). *Ann Rheum Dis* 2016 Aug; 75 (8):1438–1443. <https://doi.org/10.1136/annrheumdis-2015-207897> PMID: 26242443

17. Arends S, Brouwer E, van der Veer E, Groen H, Leijnsma MK, Houtman PM, et al. Baseline predictors of response and discontinuation of tumor necrosis factor-alpha blocking therapy in ankylosing spondylitis: a prospective longitudinal observational cohort study. *Arthritis Res Ther* 2011 Jun 20; 13(3):R94. <https://doi.org/10.1186/ar3369> PMID: 21689401
18. Dougados M, Braun J, Szanto S, Combe B, Geher P, Leblanc V, et al. Nonsteroidal antiinflammatory drug intake according to the Assessment of SpondyloArthritis International Society Score in clinical trials evaluating tumor necrosis factor blockers: example of etanercept in advanced ankylosing spondylitis. *Arthritis Care Res (Hoboken)* 2012 Feb; 64(2):290–294.
19. Garrett S, Jenkinson T, Kennedy LG, Whitelock H, Gaisford P, Calin A. A new approach to defining disease status in ankylosing spondylitis: the Bath Ankylosing Spondylitis Disease Activity Index. *J Rheumatol* 1994 Dec; 21(12):2286–2291. PMID: 7699630
20. Zochling J. Measures of symptoms and disease status in ankylosing spondylitis: Ankylosing Spondylitis Disease Activity Score (ASDAS), Ankylosing Spondylitis Quality of Life Scale (ASQoL), Bath Ankylosing Spondylitis Disease Activity Index (BASDAI), Bath Ankylosing Spondylitis Functional Index (BASFI), Bath Ankylosing Spondylitis Global Score (BAS-G), Bath Ankylosing Spondylitis Metrology Index (BASMI), Dougados Functional Index (DFI), and Health Assessment Questionnaire for the Spondylarthropathies (HAQ-S). *Arthritis Care Res (Hoboken)* 2011 Nov;63 Suppl 11:S47–58.
21. Dougados M, Simon P, Braun J, Burgos-Vargas R, Maksymowych WP, Sieper J, et al. ASAS recommendations for collecting, analysing and reporting NSAID intake in clinical trials/epidemiological studies in axial spondyloarthritis. *Ann Rheum Dis* 2011 Feb; 70(2):249–251. <https://doi.org/10.1136/ard.2010.133488> PMID: 20829199
22. van der Heijde D, Sieper J, Maksymowych WP, Dougados M, Burgos-Vargas R, Landewe R, et al. 2010 Update of the international ASAS recommendations for the use of anti-TNF agents in patients with axial spondyloarthritis. *Ann Rheum Dis* 2011 Jun; 70(6):905–908. <https://doi.org/10.1136/ard.2011.151563> PMID: 21540200